

## Assuring retention of polar compounds in an Aqueous Normal Phase ANP gradient in positive ion mode LCMS - Tips & Suggestions

**When using a Cogent TYPE-C™ column such as the Diamond Hydride prepare your solvents for positive ionization in LC-MS:**

- Solvent A: 50% methanol / 50% DI water + 0.1% formic acid
- Solvent B: 90% acetonitrile / 10% DI water + 0.1% formic acid.

Aqueous Normal Phase ANP gradients need to start between 100% - 90% of Solvent B (*to retain polar compounds*) and then reduce the percent of Solvent B as much as 20% solvent B (*when 50% methanol / 50% water is used as Solvent A*).

One important advantage to note about the Cogent TYPE-C columns is that they equilibrate between gradient runs very fast (*5 minutes*) compared to HILIC columns which require up to 20-30 minutes or more.



Printed from the Chrom Resource Center  
Copyright 2025, All Rights Apply  
**MicroSolv Technology Corporation**  
9158 Industrial Blvd. NE, Leland, NC 28451

Tel: (732) 380-8900  
Fax: (910) 769-9435  
Email: [customers@mtc-usa.com](mailto:customers@mtc-usa.com)  
Website: [www.mtc-usa.com](http://www.mtc-usa.com)