



# Phenolic Compounds Separated by LCMS - AppNote


# **Commercial Rice Extract Analyzed Easily.**

Click <u>HERE</u> for Column Ordering Information.

A commercial rice extract was analyzed and peaks were assigned based on retention times and m/z values for the compounds of interest using the Cogent Phenyl Hydride Column . After method validation the developed procedure can be used to evaluate the quality of rice and to develop the best extraction procedure.







#### Peaks:

1. 3,4-Hydroxybenzoic Acid 153 m/z [M-H]-

2. Vanillic Acid 167 m/z [M-H]-

3. p-Coumaric Acid 163 m/z [M-H]-

4. Ferulic Acid 193 m/z [M-H]-

5. 3,5-Dimethoxy-4-Hydroxycinnamic Acid 223 m/z [M-H]-

## **Method Conditions**

Column: Cogent Phenyl Hydride™, 4µm, 1 00Å Catalog No.: <u>69020-05P-2</u> Dimensions: 2.1 x 50 mm Mobile Phase: A: DI Water / 0.1% Formic Acid (v/v)

B: Acetonitrile / 0.1% Formic Acid (v/v)

### Gradient:

Time (minutes)	%B	
0	10	
5	20	

6	20	
7	10	
Post Tim	ne: 3 minutes	
Injection	ι <b>vol.:</b> 1μL	
Flow rate	e: 0.4 mL/minute	
Detectio	n: ESI – NEG - PerkinElm	er Flexar SQ 300 Mass Spectrometer
Samples	):	
Figure /	A: Commercial rice extract	was analyzed.
Figure I	B: Peak of 3,5-Dimethoxy-	4-Hydroxycinnamic Acid in commercial rice extract overlaid witl
the		
peak fo	r 25 ppm standard.	
<b>t</b> o: 0.4 r	ninutes	

**Note:** Rice is a staple food in many countries. It contains phenolic compounds which have anticancer, antioxidant, and antimutagenic effects. It is important to analyze rice extracts to confirm the content of the phenolic compounds in rice.



### Attachment

No 297 Phenolic Compounds Separated pdf 0.3 Mb Download File

Printed from the Chrom Resource Center Copyright 2025, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451 Tel: (732) 380-8900 Fax: (910) 769-9435 Email: customers@mtc-usa.com Website: www.mtc-usa.com