


1,4-Benzodiazepines in Urine Analyzed with ANP - AppNote

1,4-Benzodiazepine compounds from Urine Samples were analyzed successfully after Solid Phase Extraction (SPE).

Four available compounds (shown below) were well retained and separated. The procedure could be used for determination of this class of compounds in urine samples and other body fluids.



Peaks:

- 1. Temazepam 301.0739 m/z [M+H]+
 - 2. Diazepam 285.0790 [M+H]+
- 3. Nordiazepam 271.0633 [M+H]+
- 4. Midazolam 326.0855 [M+H]+

Method Conditions

Column: Cogent Diamond Hydride 2.0™, 2.2µm, 120Å

Catalog No.: 70200-05P-2

Dimensions: 2.1 x 50mm

Solvents:

A: DI H20 / 0 .1% formic Acid (v/v)
B: Acetonitrile/ 0.1% formic Acid (v/v)

Gradient:

time	%B
0	85
6	70
7	20
9	20
10	85

Injection vol.: 1µL

Flow rate: 0.4 ml / minute

Detection: ESI – POS – Perkin Elmer AxION 2 TOF Mass Spectrometer

Samples:

Extraction method: Spiked urine sample was loaded into SPE cartridge I (Clean Screen Xcel™ purchased from UCT Bristol, PA, USA) and eluted with 0.78mL of acetonitrile, 200μL of 2-propanol, 20μL of ammonia.

After the elution, the sample was dried under N2 gas and dissolved in $100\mu L$ of 50:50 methanol / DI H2O / 0.1% formic acid.

NOTE: Before injection, the 10 ppm spiked sample was filtered through a 0 .45µm AQ™ Brand Nylon Syringe Filter (MicroSolv Tech Corp).



Attachments

No 305 1,4-Benzodiazepines in Urine.pdf 0.3 Mb Download File

Printed from the Chrom Resource Center Copyright 2025, All Rights Apply **MicroSolv Technology Corporation** 9158 Industrial Blvd. NE, Leland, NC 28451

Fax: (910) 769-9435 Email: customers@mtc-usa.com

Website: www.mtc-usa.com

Tel: (732) 380-8900